

THE $\Lambda\Lambda - \Xi N$ COUPLING AND THE BINDING ENERGY OF ${}^6_{\Lambda\Lambda}\text{He}$

I. R. AFNAN

*Department of Physics, The Flinders University of South Australia
G.P.O. Box 2100, Adelaide 5001, Australia*

To understand the difference between the value of the nn and $\Lambda\Lambda$ two-body matrix elements, we examine the role of the coupling between the $\Lambda\Lambda$ and ΞN channels, first within the framework of $SU(3)$, and second in the numerical results for the binding energy of ${}^6_{\Lambda\Lambda}\text{He}$. We find that it is essential to include the coupled channel nature of the BB interaction as we proceed from strangeness zero to the strangeness -1 and -2 channels if we are to understand the relative magnitude of the two-body matrix elements in the different strangeness channels.

1 Introduction

The interest in the strangeness $S = -2$ baryon-baryon (BB) interaction was initially motivated by Jaffe's¹ quark model prediction of the existence of a flavor singlet $[uuddss]$ state, known as the H particle. More recently, the interest in the role of strangeness in neutron stars, and the possible production of nuclei with strangeness $S \leq -2$ at RHIC, has given further impetus to extend the BB interaction to the $S \leq -2$ channel.

At present the only experimental data on the $\Lambda\Lambda$ interaction are the three observed $\Lambda\Lambda$ hypernuclei ${}^6_{\Lambda\Lambda}\text{He}$,² ${}^{10}_{\Lambda\Lambda}\text{Be}$,^{3,4} and ${}^{13}_{\Lambda\Lambda}\text{B}$.^{5,6} These invariably give effective S-wave matrix elements of

$$\begin{aligned}\Delta B_{\Lambda\Lambda}({}^6_{\Lambda\Lambda}\text{He}) &= 10.9 \pm 0.5 - 6.14 \pm 0.05 \approx 4.7 \text{ MeV} \\ \Delta B_{\Lambda\Lambda}({}^{10}_{\Lambda\Lambda}\text{Be}) &= 17.7 \pm 0.4 - 13.42 \pm 0.08 \approx 4.3 \text{ MeV} \\ \Delta B_{\Lambda\Lambda}({}^{13}_{\Lambda\Lambda}\text{B}) &= 27.5 \pm 0.7 - 11.69 \pm 0.12 \approx 4.8 \text{ MeV},\end{aligned}\tag{1}$$

which suggests that the matrix element of the $\Lambda\Lambda$ interaction in light nuclei is $-\langle V_{\Lambda\Lambda} \rangle \approx 4 - 5$ MeV. This value is to be compared with the corresponding nn matrix element $-\langle V_{nn} \rangle \approx 6 - 7$ MeV, and the ΛN matrix element $-\langle V_{\Lambda N} \rangle \approx 2 - 3$ MeV. As a result we have that $-\langle V_{nn} \rangle > -\langle V_{\Lambda\Lambda} \rangle > -\langle V_{\Lambda N} \rangle$. The question we would like to address in the present report is: Can we understand this relative magnitude of the $S = 0, -1, -2$ matrix elements in light nuclei in terms of flavor $SU(3)$ at the meson-baryon level as given in a one-boson-exchange (OBE) potential? Or do we need to include explicit quark-gluon degrees of freedom? We will demonstrate that the relative magnitude of the matrix elements in the $S = 0, -1, -2$ can be understood only if we include the coupling between the ΛN and ΣN in the $S = -1$ channels, and the $\Lambda\Lambda$ and ΞN in the $S = -2$ channel.

2 The Coupled Channel Problem in Flavor $SU(3)$

Let us assume that flavor $SU(3)$ is a perfect symmetry of strong interaction, then the baryon octet ($B = N, \Lambda, \Sigma, \Xi$) forms a degenerate multiplet, and the BB system will be in one of the six irreducible representations of $\{8\} \otimes \{8\}$, i.e.

$$\{8\} \otimes \{8\} = [\{1\} \oplus \{8_s\} \oplus \{27\}] \oplus [\{8_a\} \oplus \{10\} \oplus \{\bar{10}\}],\tag{2}$$

where $\{8_a\}$, $\{10\}$ and $\{\bar{1}0\}$ are the flavor anti-symmetric channels, while the $\{1\}$, $\{8_s\}$ and $\{27\}$ are the flavor symmetric channels. The flavor wave function in the $^1S_0$ partial wave for the nn , ΛN and $\Lambda\Lambda$ channels can now be written with the help of the $SU(3)$ Clebsch-Gordan coefficients,⁷ as

$$\begin{aligned} |nn\rangle &= |\{27\}\rangle \\ |\Lambda N\rangle &= \sqrt{\frac{36}{40}} |\{27\}\rangle - \sqrt{\frac{4}{40}} |\{8_s\}\rangle \\ |\Lambda\Lambda\rangle &= \sqrt{\frac{27}{40}} |\{27\}\rangle - \sqrt{\frac{8}{40}} |\{8_s\}\rangle - \sqrt{\frac{5}{40}} |\{1\}\rangle . \end{aligned} \quad (3)$$

Here the $\{27\}$ representation is the only component of the nn flavor wave function. Since the nn interaction in the $\{27\}$ representation is strongly attractive to the extent that it almost supports a bound state, and both the ΛN and $\Lambda\Lambda$ wave flavor wave functions have their largest component in the $\{27\}$ representation, we expect the ΛN and $\Lambda\Lambda$ interactions in the $^1S_0$ to also be dominated by the $\{27\}$ representation. As a result the interactions in the $S = 0, -1$ and -2 are expected to satisfy the relation

$$-\langle V_{nn} \rangle > -\langle V_{\Lambda N} \rangle > -\langle V_{\Lambda\Lambda} \rangle . \quad (4)$$

This is in contradiction with the experimental observation from the three $\Lambda\Lambda$ hypernuclei which requires that $-\langle V_{\Lambda\Lambda} \rangle > -\langle V_{\Lambda N} \rangle$. The inclusion of the $\{8_s\}$ and $\{1\}$ component of the flavor wave function into our analysis may not improve the situation since there are indications from OBE potentials that the interaction in these two channels are repulsive⁸ which would reduce the $\Lambda\Lambda$ matrix element more than the ΛN matrix element.

Since flavor $SU(3)$ is broken in nature to the extent that the masses of the N , Λ , Σ and Ξ are not equal, the BB problem in the $S = -1, -2$ reduces to a coupled channel problem with two or more thresholds in each strangeness channel. Thus for $S = -1$ we have the ΛN and ΣN thresholds which are separated by some 80 MeV; for the $S = -2$ we have the $\Lambda\Lambda$, ΞN and $\Sigma\Sigma$ thresholds with the separation between the $\Lambda\Lambda$ and ΞN channels being ≈ 25 MeV. This suggests that the coupling in the $S = -2$ may be more important than is the case in the $S = -1$ channel. To examine the effect of the baryon mass splitting on ΛN and $\Lambda\Lambda$ matrix elements, we introduce effective potentials that include the coupled channel effects. These potentials, to lowest order in the coupling, are given by

$$\begin{aligned} \langle V_{\Lambda N}^{\text{eff}} \rangle &\approx \langle V_{\Lambda N} \rangle - \frac{|\langle \Lambda N | V | \Sigma N \rangle|^2}{\Delta E_{YN}} ; & \Delta E_{YN} &\approx 80 \text{ MeV} \\ \langle V_{\Lambda\Lambda}^{\text{eff}} \rangle &\approx \langle V_{\Lambda\Lambda} \rangle - \frac{|\langle \Lambda\Lambda | V | N\Xi \rangle|^2}{\Delta E_{\Lambda\Lambda}} ; & \Delta E_{\Lambda\Lambda} &\approx 25 \text{ MeV} . \end{aligned} \quad (5)$$

The fact that the thresholds in the $S = -2$ channel are closer than in the $S = -1$ channel suggests that we get more attraction in the $S = -2$ effective interaction due to the coupled channel nature of the problem, than in the $S = -1$ channel. If in addition we assume that the matrix elements that determine the strength of the

coupling can be extracted from $SU(3)$, i.e. flavor $SU(3)$ is approximately a good symmetry, then

$$\begin{aligned}\langle \Lambda\Lambda|V|\Xi N\rangle &= -\frac{18}{40}V_{\{27\}} + \frac{8}{40}V_{\{8_s\}} + \frac{10}{40}V_{\{1\}} \\ \langle \Lambda N|V|\Sigma N\rangle &= -\frac{12}{40}V_{\{27\}} + \frac{3}{40}V_{\{8_s\}} .\end{aligned}\tag{6}$$

If the interaction in the $\{27\}$ representation is dominant, then the coupling between the $\Lambda\Lambda$ and the ΞN channels is stronger than is the case between the ΛN and the ΣN channels. This again supports the argument that the coupling resulting from the mass splitting gives more attraction in the $\Lambda\Lambda$ channel than in the ΛN channel. Thus even though the $^1S_0$ interaction in the $\Lambda\Lambda$ channel may be weak (small scattering length), the additional attraction resulting from the coupling due to the baryon mass splitting could result in a scattering length that is comparable to the nn scattering length as first suggested by Dover.⁹

Let us now turn to the $\Lambda\Lambda$ interaction having established that the coupling between the $\Lambda\Lambda$ and ΞN channels is more important than is the case in the $S = -1$ channel. Since the experimental data put a constraint on the scattering amplitude that included the additional attraction due to the coupled channel nature of the problem, the $\Lambda\Lambda$ potential in free space, in the absence of coupling, is considerably stronger than is the case when the coupling is included. However in the medium, the coupling is suppressed as a result of the fact that $\Lambda + \Lambda \rightarrow \Xi + N$ is Pauli blocked. From this we may conclude that while in free space

$$\langle V_{\Lambda\Lambda}^{\text{eff}} \rangle_{\text{with coupling}} = \langle V_{\Lambda\Lambda}^{\text{eff}} \rangle_{\text{no coupling}} ,$$

in the medium

$$\langle V_{\Lambda\Lambda}^{\text{eff}} \rangle_{\text{with coupling}} < \langle V_{\Lambda\Lambda}^{\text{eff}} \rangle_{\text{no coupling}} .$$

In other words, if the $\Lambda\Lambda$ amplitude in free space is attractive enough to almost support a bound state, in the medium the Pauli principle suppresses this amplitude so that the effective matrix element of the $\Lambda\Lambda$ interaction is smaller than is the case for the nn interaction.

The above qualitative analysis is based on the assumption that flavor $SU(3)$ gives a good estimate of the relative magnitude of matrix elements and the breaking of $SU(3)$ is only in the masses of the baryons. To see if these arguments are valid, in the next section we will report on the construction of a potential in the $S = -2$ channel based on the $SU(3)$ rotation of the Nijmegen model D potential,¹⁰ and then proceed in Sec. 4 to use the resulting potential in the calculation of the binding energy of ${}^6_{\Lambda\Lambda}\text{He}$. Finally, in Sec. 5 we will present some concluding remarks.

3 The $S = -2$ BB Potentials

To perform an $SU(3)$ rotation on an OBE potential defined in $S = 0, -1$ channels, we need to write the Lagrangian in terms of the baryon octet with the mesons as

either a singlet or a member of an octet. If the interaction is taken to be of the Yukawa type, the Lagrangian takes the form

$$\mathcal{L}_{int} = - \{ g_{\{1\}} [B^\dagger B]_{\{1\}} M_{\{1\}} + g_{\{8_s\}} [B^\dagger B]_{\{8_s\}} M_{\{8_s\}} + g_{\{8_a\}} [B^\dagger B]_{\{8_a\}} M_{\{8_a\}} \} , \quad (7)$$

where B and M are the field operators for the baryon and mesons. In writing the above Lagrangian, which is a scalar, we have coupled the initial and final baryons to a flavor singlet or an octet. Since there are two irreducible octet representations, we need a different coupling constant for each of the representations. This Lagrangian has one coupling constant for each singlet meson $g_{\{1\}}$, and two coupling constants $g_{\{8_s\}}$ and $g_{\{8_a\}}$ for each meson octet. These coupling constants can then be determined by fitting the experimental data.

The Nijmegen potential¹⁰ model D takes for the exchange mesons the pseudoscalar octet $\{\pi, \eta, \eta', K\}$, the vector octet $\{\rho, \phi, \omega, K^*\}$ and a scalar meson $\{\epsilon\}$. The masses of the mesons and baryons are taken from experiment, while the coupling constants are adjusted to fit the data in the $S = 0, -1$. This, in principle, determines the long range part of the potential which should be described in terms of meson-baryon degrees of freedom. These same coupling constants can be used to construct the OBE potential for $S \leq -2$. For the short range part of the interaction, model D assumes a hard core, which is an additional parameter that could be different in the $S = 0$ and $S = -1$ channels. The breaking of flavor $SU(3)$ is due to the use of physical masses for the baryons and mesons, and the choice of different hard core radii in the $S = 0, -1$.

We expect the short range part of the BB interaction to be expressed in terms of quark-gluon degrees of freedom. In a valence quark model description of this short range interaction the number of strange quarks increases as we go from $S = 0$ to $S = -1$ and then $S \leq -2$. As the number of strange quarks increases, we expect the role of the Pauli principle at the quark level to change. As a result the short range part of the BB interaction should be S dependent, and cannot be determined from the $S = 0, 1$ channels at the meson-baryon level in terms of a flavor $SU(3)$ rotation of the OBE potential. Thus to determine the BB interaction in the $S = -2$ channel, we need to introduce at least one additional parameter that determines the short range interaction.

This procedure is followed by Carr *et al.*¹¹ They only considered S -wave interaction and ignored the tensor interaction. Their potential for the exchange of the i^{th} meson is of the form

$$V_i(r) = V_c^{(i)}(r) + \vec{\sigma}_1 \cdot \vec{\sigma}_2 V_\sigma^{(i)}(r) , \quad (8)$$

where the radial potential $V_\alpha^{(i)}$, $\alpha = c, \sigma$ for a meson of mass m_i is taken to be

$$V_\alpha^{(i)}(r) = V_0^{(i)} \left[\frac{e^{-m_i r}}{m_i r} - C \left(\frac{M}{m_i} \right) \frac{e^{-Mr}}{Mr} \right] ; \quad \alpha = c, \sigma . \quad (9)$$

To guarantee a one parameter short range repulsion, a mass $M = 2500$ MeV is chosen in all partial waves. Then the parameter C determined the strength of the short range interaction. This new parameter C is constrained to ensure that the potential for $r \geq 0.7$ fm is unchanged and the short range interaction is always repulsive. In Fig. 1 we present the $\Lambda\Lambda$ potential $V_{\Lambda\Lambda}$ and the ΞN potential $V_{\Xi N}$.




Figure 1. The $S = -2$, $^1S_0$ one-boson-exchange potential $V_{\Lambda\Lambda}$ and $V_{\Xi N}$.

In addition to the OBE potential (solid line), the figure includes three potentials in which the parameter C is adjusted to support a bound state (C), generate an anti-bound state (B), or have no bound state at all (A).

To examine the role of the coupling between the $\Lambda\Lambda$ and ΞN channels in ${}_{\Lambda\Lambda}^6\text{He}$, we need to include this coupling exactly. This can be achieved most simply within a model in which ${}_{\Lambda\Lambda}^6\text{He}$ is taken to be a $\Lambda\Lambda\alpha$ three-body system described by the Faddeev equations. These equations can be reduced to coupled one dimensional integral equations if the interactions are assumed to be separable. For this Carr *et al*¹¹ generated a set of separable potentials that give the same effective range parameters as meson exchange potentials A, B and C. These are referred to as SA, SB and SC. The effective range parameters for both the local and the separable potentials are given in Table 1. Here we observe that the potentials B and SB have a scattering length that is comparable to the nn scattering length.

Table 1. The effective range parameters and binding energy of the two-body $\Lambda\Lambda - \Xi N$ system for the local potentials A, B and C and the equivalent separable potentials SA, SB and SC.

Pot.	$a_{\Lambda\Lambda}$	$r_{\Lambda\Lambda}$	$a_{\Xi N}$	$r_{\Xi N}$	B.E. (MeV)
A	-1.91	3.36	-2.12-0.75i	3.45-0.45i	UB
SA	-1.90	3.33	-2.08-0.81i	3.44-0.22i	UB
B	-21.1	1.86	-2.05-6.53i	2.12-0.21i	UB
SB	-21.0	2.54	-2.07-6.52i	2.62-0.15i	UB
C	7.82	1.41	3.08-5.26i	1.74-0.144i	0.71
SC	7.84	1.48	3.05-5.28i	1.45+0.074i	0.71

4 Binding Energy ${}_{\Lambda\Lambda}{}^6\text{He}$

In reducing the six body problem of ${}_{\Lambda\Lambda}{}^6\text{He}$ to a three-body problem we gain the advantage of being able to solve the problem exactly using the Faddeev equations. In particular, the treatment of the coupled channel $\Lambda\Lambda - \Xi N$ is included in full. The main disadvantage of this three-body model is the limitation in the handling of the Pauli principle. This is most pronounced when the two Λ s convert to ΞN , as the resulting nucleon should be in an antisymmetric state with respect to the nucleons in the α particle. Since the latter is treated as an elementary particle with no internal degrees of freedom, it is not possible to guarantee that the Pauli principle is satisfied. This effect is most pronounced when the nucleon and α particle are in relative S -wave. There are two ways of overcoming this problem in an approximate way. The first is to assume that the S -wave interaction supports a bound state in the absence of the Pauli exclusion principle. We then can remove this bound state from the spectrum of the Hamiltonian without any modification to the phase shifts. The resultant amplitude has the Pauli forbidden state removed. The second approach is to take the S -wave $N - \alpha$ interaction to be repulsive. The two methods give almost identical results for the binding energy of ${}^6\text{Li}$.¹² Carr *et al*¹¹ used the latter approximation as it results in a rank one potential and therefore a simpler set of Faddeev equations for the three-body model of ${}_{\Lambda\Lambda}{}^6\text{He}$.

The second complication in turning ${}_{\Lambda\Lambda}{}^6\text{He}$ into a three-body problem is the need to know the $\Lambda - \alpha$ and $\Xi - \alpha$ interactions. These interactions should be consistent with the two-body BB interaction. Here again, Carr *et al*¹¹ resorted to simple models for these interactions. For the case of the $\Lambda - \alpha$ interaction, they made use of the G -matrix parametrization¹³ of the Nijmegen model D potential to fold this interaction with ${}^4\text{He}$ density, while for the $\Xi - \alpha$ interaction they employed a Wood-Saxon optical potential.

Finally, since the conversion of the $\Lambda\Lambda$ to ΞN releases ≈ 25 MeV of energy, that energy could go into the excitation of the α particle. Although it is possible to include such excitation by assuming the α particle can be in one of two states with mass separation comparable to the energy of the first excited state of ${}^4\text{He}$, Carr *et al*¹¹ chose not to include this excitation as the effective coupling to the $\Xi N\alpha$ channel is suppressed because the S -wave αN interaction has been taken to be repulsive, i.e. because of Pauli blocking.

To examine the relative importance of the coupling between the $\Lambda\Lambda$ and ΞN in free space, and in the nuclear medium where the Pauli blocking plays an important role, we consider three models for the binding energy of ${}_{\Lambda\Lambda}{}^6\text{He}$. These are:

1. In model 1 we include the full coupling between the $\Lambda\Lambda$ and the ΞN . This reduces the three-body model for ${}_{\Lambda\Lambda}{}^6\text{He}$ to the three-body coupled channel problem $\Lambda\Lambda\alpha - \Xi N\alpha$, which can be solved exactly to give the binding energy for the three-body system under consideration. This is an exact solution to the problem within the limitations stated above. Here we note that this system is not a pure $\Lambda\Lambda\alpha$ system, but has a certain probability of being in a $\Xi N\alpha$ state.
2. In model 2 we include the coupling between the $\Lambda\Lambda$ and ΞN channels in calculating the two-body T -matrix that is the input to the Faddeev equations,

but carry out the three-body calculation within the $\Lambda\Lambda\alpha$ space only. This is equivalent to setting $T_{\Lambda\Xi} = 0$, which gives us a measure of the importance of the coupling between the $\Lambda\Lambda$ and ΞN in nuclear matter, and the effect of the Pauli blocking. In this case the probability of finding the system in $\Xi N\alpha$ state is zero.

3. Finally, in model 3 we assume that the coupling between the $\Lambda\Lambda$ and ΞN can be neglected at the two-body level in free space, i.e. $V_{\Lambda\Xi} = 0$. In this case we construct a two-body $\Lambda\Lambda$ interaction that gives the same $\Lambda\Lambda$ effective range parameters as the coupled channel problem. Thus for model B, the $\Lambda\Lambda$ potential, in the absence of coupling, has an anti-bound state and a scattering length comparable to the nn scattering length.

In Table 2 we present the results of Carr *et al*¹¹ for the above three models using the potentials SA, SB and SC for the $\Lambda\Lambda - \Xi N$ interaction. We observe that:

- The experimental result of 10.9 ± 0.6 lies between the “exact” results (model 1) for potentials SA and SB, and closer to SB. This suggests that within the present model the short range parameter C should be such that the $\Lambda\Lambda$ scattering length is large and comparable to the nn scattering length if we are to reproduce the binding energy of ${}_{\Lambda\Lambda}^6\text{He}$. Therefore we can use the binding energy of one of the $\Lambda\Lambda$ hypernuclei to fix the short range part of the interaction in the $S = -2$ channel and then proceed to use the resulting potential for other $S = -2$ hypernuclei.
- If we now compare the results of model 1 and model 2 in Table 2, we observe that the contribution of $T_{\Lambda\Xi}$ is small (≈ 0.5 MeV) if our potential would give a binding energy comparable to the experimental result. This relatively small reduction in binding energy when the $\Lambda\Lambda - \Xi N$ coupling is included is a measure of the importance of Pauli blocking, even in a nucleus as light as ${}_{\Lambda\Lambda}^6\text{He}$. This

Table 2. The binding energy of ${}_{\Lambda\Lambda}^6\text{He}$ for the three models under consideration and for the three interactions SA, SB and SC.

	$a_{\Lambda\Lambda}$ (fm)	BE(model 1) ($\Lambda\Lambda\alpha - \Xi N\alpha$) (MeV)	BE(model 2) ($\Lambda\Lambda\alpha; T_{\Lambda\Xi} = 0$) (MeV)	BE(model 3) ($\Lambda\Lambda\alpha; V_{\Lambda\Xi} = 0$) (MeV)
SA	-1.91	9.738	9.5078	10.007
SB	-21.1	12.268	11.606	14.134
SC	7.82	15.912	14.533	17.842
Exp.		10.9 ± 0.6		

small change in binding energy due to coupling in the nuclear medium does in part justify the exclusion of the excitation of the α particle in the present models.

- On the other hand, if we compare the results of model 1 and model 3 in Table 2, we may conclude that turning off the coupling between the $\Lambda\Lambda$ and ΞN channels at the two-body level, in free space, is in fact a very poor approximation. The difference of ≈ 2 MeV between model 1 and model 3 is a substantial fraction of the value of the $\Lambda\Lambda$ matrix element (a change of $\approx 30\%$). This difference is, to lowest order, the contribution of the coupling between the $\Lambda\Lambda$ and ΞN to the attraction in the effective $\Lambda\Lambda$ matrix element. To see this, we write the binding energy of ${}^6_{\Lambda\Lambda}\text{He}$ as the matrix element of the Hamiltonian, i.e.

$$-B.E. = \langle H \rangle = K.E. + 2\langle V_{\Lambda\alpha} \rangle + \langle V_{\Lambda\Lambda}^{\text{eff}} \rangle, \quad (10)$$

where $\langle V_{\Lambda\Lambda}^{\text{eff}} \rangle$ is the effective $\Lambda\Lambda$ matrix elements in the medium. If there is no coupling between the $\Lambda\Lambda$ and ΞN channels in free space then

$$\langle V_{\Lambda\Lambda}^{\text{eff}} \rangle = \langle V_{\Lambda\Lambda} \rangle_{\text{no coupling}} \quad \text{for} \quad V_{\Lambda\Xi} = 0. \quad (11)$$

On the other hand, the inclusion of the coupling between the $\Lambda\Lambda$ and the ΞN in free space reduces the strength of the $\Lambda\Lambda$ potential, $V_{\Lambda\Lambda}$, so that the $\Lambda\Lambda$ potential and the additional attraction due to the coupling results in an effective matrix element that is fixed by experiment. If we assume that the Pauli blocking removes the contribution of the coupling between the $\Lambda\Lambda$ and ΞN in the nuclear medium, i.e. the difference between the result of model 1 and model 2 in Table 2 is zero, then

$$\langle V_{\Lambda\Lambda}^{\text{eff}} \rangle = \langle V_{\Lambda\Lambda} \rangle_{\text{with coupling}} \quad \text{for} \quad V_{\Lambda\Xi} \neq 0. \quad (12)$$

But from Eq. (10) we know that the difference between the binding energy with and without coupling between the $\Lambda\Lambda$ and the ΞN channels in free space is the contribution of the coupling to the attraction, i.e.

$$\begin{aligned} BE(\text{model 1}) - BE(\text{model 3}) &\approx -\langle V_{\Lambda\Lambda} \rangle_{\text{with coupling}} + \langle V_{\Lambda\Lambda} \rangle_{\text{no coupling}} \\ &\approx \frac{|\langle \Lambda\Lambda | V | N\Xi \rangle|^2}{\Delta E_{\Lambda\Lambda}} \approx -2 \text{ MeV}. \end{aligned} \quad (13)$$

This result is approximately the difference between the nn and $\Lambda\Lambda$ matrix elements in light nuclei.

This analysis basically establishes the fact that it is essential to include the coupling between the $\Lambda\Lambda$ and ΞN channels in free space even though the coupling is suppressed in the medium due to Pauli blocking.

5 Conclusion

From the above analysis based on the validity of flavor $SU(3)$ for matrix elements of the potential, we can understand how the coupling resulting from mass

differences between the members of the baryon octet could justify the fact that $-\langle V_{\Lambda\Lambda} \rangle > -\langle V_{\Lambda N} \rangle$. In fact flavor $SU(3)$ suggests that the coupling in the $S = -2$ is substantially more important than is the case in the $S = -1$ channel. This is a result of the fact that: (i) The $\Lambda\Lambda$ threshold and the ΞN threshold are closer than the ΛN and ΣN thresholds. (ii) The coupling matrix element for the transition from $\Lambda\Lambda$ to ΞN is larger than the coupling matrix elements for the transition from ΛN to ΣN .

With the help of flavor $SU(3)$ rotation we showed how we can construct the long range part of the potential based on OBE for $S \leq -2$ given that the meson baryon coupling constants are fixed by the data in the $S = 0, -1$ channels. However, this flavor $SU(3)$ rotation does not determine the short range part of the interaction. In fact valence quark models for the BB system would give an S dependence to the short range data, and in an OBE model this requires additional data, e.g. the binding energy of $\Lambda\Lambda$ hypernuclei.

Using this $SU(3)$ rotation Carr *et al*¹¹ constructed an extension of the Nijmegen model D potential to the $S = -2$ channel with a soft cut-off that does not change the potential outside of 0.7 fm. Finally, Carr *et al*¹¹ were able to show that a $\Lambda\Lambda$ scattering length comparable to the nn scattering length gives approximately the correct binding energy for ${}_{\Lambda\Lambda}^6\text{He}$. This suggests that the experimental binding energy of ${}_{\Lambda\Lambda}^6\text{He}$ is consistent with the strength of the BB potential as predicted by flavor $SU(3)$ rotation of the OBE potentials. More important it confirms the prediction of Dover⁹ that the $\Lambda\Lambda$ scattering length is comparable to the nn scattering length in the ${}^1S_0$.

With the inclusion of coupling between the $\Lambda\Lambda$ and ΞN channel there is no need to include repulsive three-body forces to calculate the correct binding energy for ${}_{\Lambda\Lambda}^6\text{He}$. This is a result of the fact that the Pauli blocking of the process $\Lambda\Lambda \rightarrow \Xi N$ in the nuclear medium reduces the effective attraction in the $\Lambda\Lambda$ channel as compared to the attraction in free space. As a result the $\Lambda\Lambda$ matrix element in light nuclei is less than the nn matrix element by about 2 MeV, which is consistent with the experimental results from $\Lambda\Lambda$ hypernuclei. This is despite the fact that the scattering length for the $\Lambda\Lambda$ is comparable to the nn scattering length.

Finally, we can understand the result that

$$-\langle V_{nn} \rangle > -\langle V_{\Lambda\Lambda} \rangle > -\langle V_{\Lambda N} \rangle, \quad (14)$$

on the basis of the coupled channel nature of the BB interaction in the $S = -1, -2$ channels resulting from flavor $SU(3)$ breaking in the mass of the baryon octet.

Acknowledgments

The author would like to thank the Australian Research Council for their financial support. He is indebted to his collaborators S.B. Carr and B.F. Gibson for many enlightening discussions and helpful suggestions.

References

1. A. Jaffe, *Phys. Rev. Lett.* **38**, 195 (1977).

2. D.J. Prowse, *Phys. Rev. Lett.* **17**, 782 (1966).
3. M. Danysz *et al*, *Nucl. Phys.* **49**, 121 (1963).
4. R.H. Dalitz *et al*, *Proc. R. Soc. London A* **426**, 1 (1989).
5. S. Aoki *et al*, *Phys. Rev. Lett.* **65**, 1729 (1990); *Prog. Theo. Phys.* **85**, 1287 (1991).
6. C.B. Dover, D.J. Millener, A. Gal and D.H. Davis, *Phys. Rev., C* **44**, 1905 (1991).
7. J.J. de Swart, *Rev. Mod. Phys.* **35**, 916 (1963).
8. Th.A. Rijken, P.M.M. Maessen and J.J. de Swart, *Nucl. Phys. A* **547**, 245c (1992).
9. C.B. Dover, Properties & Interactions of Hypernuclei, eds. B.F. Gibson, P.D. Barnes and K. Nakai (World Scientific, Singapore, 1994), p 1.
10. M.M. Nagel, Th.A. Rijken and J.J. de Swart, *Phys. Rev. D* **15**, 2547 (1977).
11. S.B. Carr, I.R. Afnan and B.F. Gibson, *Nucl. Phys. A* **625**, 143 (1997).
12. D.R. Lehman, M. Rai and A. Ghovanlou *Phys. Rev. C* **17**, 744 (1978).
13. Y. Yamamoto and H. Bando, *Prog. Theo. Phys.* **73**, 905 (1985)